

Deformation quantization using groupoids. Case of toric manifolds

Frédéric CADET

February 1, 2008

Abstract

In the framework of C^* -algebraic deformation quantization we propose a notion of deformation groupoid which could apply to known examples *e.g.* Connes' tangent groupoid of a manifold, its generalisation by Landsman and Ramazan, Rieffel's noncommutative torus, and even Landi's noncommutative 4-sphere. We construct such groupoid for a wide class of $\mathbb{T}^n$ -spaces, that generalizes the one given for $\mathbb{C}^n$ by Bellissard and Vittot. In particular, using the geometric properties of the moment map discovered in the '80s by Atiyah, Delzant, Guillemin and Sternberg, it provides a C^* -algebraic deformation quantization for all toric manifolds, including the 2-sphere and all complex projective spaces.

Introduction

Quantization is going from classical mechanics to quantum mechanics. In classical hamiltonian picture, a physical system is described by its phase space, a Poisson manifold M , which Poisson bivector is denoted by Ω , and associated Poisson bracket of two observables $\mathbf{f}, \mathbf{f}' \in C^\infty(M)$ by:

$$\{\mathbf{f}, \mathbf{f}'\}_\Omega \stackrel{def}{=} \langle d\mathbf{f} \otimes d\mathbf{f}' | \Omega \rangle.$$

On the other hand a quantum system is described, in Heisenberg's picture, by replacing the classical algebra $C^\infty(M)$ of observables by an algebra of operators on a Hilbert space, in a linear and involution-preserving way. Moreover the Poisson bracket $\{\mathbf{f}, \mathbf{f}'\}_\Omega$ has to be replaced by $\frac{1}{i\hbar}[f, f']$, the commutator of f and f' , the quantum analogs of $\mathbf{f}$ and $\mathbf{f}'$, divided by $\imath = \sqrt{-1}$ and by $\hbar$, which is Planck's physical constant. Dirac postulated in the '30s that this correspondance from renormalized commutators to Poisson bracket is given by a limit process when Planck's constant goes to zero.

In the '70s the notion of *deformation quantization* was introduced in [BFFLS] as an attempt to give a precise definition to Dirac's principle. One of the ideas was to use a field of algebras $(\mathcal{A}_\hbar)$ parametrized by a real number $\hbar$, such that, for $\hbar = 0$, the algebra $\mathcal{A}_0$ is the commutative algebra of classical observables, and for $\hbar \neq 0$, $\mathcal{A}_\hbar$ is the noncommutative algebra of quantum observables. At the

beginning of the '90s Rieffel proposed in [Rie89, Rie] a topological definition of the limit process making use of the previously known notion of continuous field of C^* -algebras [DF]. The following is a new definition close Rieffel's one, but a little more general so that there is a direct adaptation to groupoid C^* -algebras.

Definition 0.1 — Let $(\mathcal{A}_h)_{h \in X}$ be a continuous field of C^* -algebras parametrized by a locally compact subset X of $\mathbb{R}$ containing 0 as a limit point; we denote $\mathcal{A} = \bigcup_{h \in X} \mathcal{A}_h$ the associated topological bundle over X . Let $\mathcal{Q}$ be a sub- $*$ -algebra of the C^* -algebra $C_0(X, \mathcal{A})$ of continuous sections of $\mathcal{A}$. We say that $((\mathcal{A}_h)_{h \in X}, \mathcal{Q})$ is a *deformation* if:

- The space $\mathcal{Q}_0 = \{f_0 \in \mathcal{A}_0 \mid f \in \mathcal{Q}\}$ is dense in $\mathcal{A}_0$.
- There is a map $\mathcal{Q}_0 \times \mathcal{Q}_0 \longrightarrow \mathcal{Q}_0$, denoted by:

$$(a, b) \longmapsto \{a, b\}_0,$$

such that, for every $f, f' \in \mathcal{Q}$, $\{f_0, f'_0\}_0$ is the continuous extension at zero of the continuous section $h \longmapsto \frac{1}{i\hbar} [f_h, f'_h]$ defined on $X - \{0\}$ i.e. :

$$\lim_{\substack{h \rightarrow 0 \\ h \neq 0}} \frac{1}{i\hbar} [f_h, f'_h] = \{f_0, f'_0\}_0.$$

Then $\mathcal{A}_0$ is a commutative C^* -algebra and $\mathcal{Q}_0$ is a sub- $*$ -algebra, and moreover $\mathcal{Q}_0$ has a structure of Poisson algebra, for the bracket $\{.,.\}_0$.

Definition 0.2 — Let (M, Ω) be a Poisson manifold. A *deformation of M* is a deformation $((\mathcal{A}_h)_{h \in X}, \mathcal{Q})$ endowed with an isomorphism of C^* -algebras :

$$C_0(M) \xrightarrow{\mathcal{I}} \mathcal{A}_0,$$

such that:

- $\mathcal{Q}_0 \subset \mathcal{I}(C_0^\infty(M))$
- for each $f, f' \in \mathcal{Q}$: $\{f_0, f'_0\}_0 = \mathcal{I}(\{\mathcal{I}^{-1}f_0, \mathcal{I}^{-1}f'_0\}_\Omega)$.

Then each linear $*$ -preserving section T of the canonical projection of $\mathcal{Q}$ onto $\mathcal{Q}_0$ gives rise to a quantization:

$$\begin{array}{ccc} \mathcal{Q}_0 & \xrightarrow{\text{quantization}} & \mathcal{Q}_h \\ & \nearrow & \nwarrow \\ & \mathcal{Q} & \end{array}$$

T

Many usual C^* -algebras can be described as C^* -algebras $C^*(G)$ of a groupoid G , using the construction given by Jean Renault in [Ren] that generalizes that of the (full and reduced) C^* -algebra(s) of a group or of a group action. A relevant

fact is that a bundle of groupoids is itself a groupoid. On the topological level Ramazan [Ram98, LR99] has established the following property:

Theorem 0.3 — *Let $(G_{\hbar})_{\hbar \in X}$ be a field of groupoids, and $G = \bigcup_{\hbar \in X} G_{\hbar}$ be the corresponding bundle which is supposed to be locally compact Hausdorff separable, having continuous Haar system, and such that the bundle map $G \xrightarrow{p} X$ is open. If G is amenable [AR] then the field $(C^*(G_{\hbar}))_{\hbar \in X}$ possesses a structure of continuous field such that the algebra of continuous sections is:*

$$C_0 \left(X, \bigcup_{\hbar \in X} C^*(G_{\hbar}) \right) = C^*(G).$$

Then it is natural to set:

Definition 0.4 — A locally compact Hausdorff separable amenable bundle of groupoids G over a locally compact subset X of $\mathbb{R}$, with continuous Haar system is a *deformation groupoid* if there exists a sub-* $\mathcal{Q}$ -algebra $\mathcal{Q}$ of sections of the continuous field $(C^*(G_{\hbar}))_{\hbar \in X}$ such that $((C^*(G_{\hbar}))_{\hbar \in X}, \mathcal{Q})$ is a deformation (resp. is a *deformation groupoid of a Poisson manifold* M if there exists $\mathcal{Q}$ and an isomorphism $\mathcal{I}$ such that $((C^*(G_{\hbar}))_{\hbar \in X}, \mathcal{Q}, \mathcal{I})$ is a deformation of M).

Finally a bundle of groupoids G over X (locally compact, Hausdorff, separable, amenable, with continuous Haar system and open projection $G \xrightarrow{p} X$) is a deformation groupoid of (M, Ω) if and only if:

1. There exists an isomorphism $C_0(M) \xrightarrow[\sim]{\mathcal{I}} C^*(G_0)$.
2. There exists a sub-* $\mathcal{Q}$ -algebra $\mathcal{Q}$ of $C^*(G)$ such that:
 - (a) $\mathcal{Q}_0$ is dense in $C^*(G_0)$,
 - (b) $\mathcal{I}^{-1}(\mathcal{Q}_0) \subset C_0^\infty(M)$,
 - (c) $\forall f, f' \in \mathcal{Q}, \lim_{\substack{\hbar \rightarrow 0 \\ \hbar \neq 0}} \frac{1}{i\hbar} [f_{\hbar}, f'_{\hbar}] = \mathcal{I}(\{ \mathcal{I}^{-1} f_0, \mathcal{I}^{-1} f'_0 \}_{\Omega})$.

Moreover we will use:

Proposition 0.5 — *Let M be a manifold, without a priori Poisson structure. If there exists a bundle of groupoids G over X (locally compact, Hausdorff, separable, amenable, with continuous Haar system) which satisfies the previous conditions 1,2a,2b, and such that*

- 2c' *for every $f, f' \in \mathcal{Q}$ the section $\hbar \longrightarrow \frac{1}{i\hbar} [f_{\hbar}, f'_{\hbar}]$ has a unique continuous extension at $\hbar = 0$ in $\mathcal{Q}_0$, which only depends on the values f_0 and f'_0 ,*

then M admits a Poisson bracket such that G is a deformation groupoid of M .

The first *example* of deformation groupoid was given by Connes in [Co] (cf. [Var01] too): the tangent groupoid of a manifold N is a deformation groupoid of the cotangent bundle T^*N endowed with its canonical symplectic structure. This was generalized by Landsman and Ramazan [LR99, Ram98, Lan98] to integrable Lie-Poisson manifolds (*i.e.* manifolds which are the dual of an integrable

Lie algebroid, endowed with the canonical Poisson structure described in [DS]. In 1990 Bellissard and Vittot [BV] constructed a deformation groupoid of $\mathbb{C}^n$ with its canonical symplectic structure, which was not a Lie groupoid. In the present paper, we generalize this construction to other $\mathbb{T}^n$ -spaces M ($\mathbb{T}^n$ denotes the n -dimensional torus); it applies to the 2-sphere and, using Delzant's results [Del88], more generally to all toric manifolds, including all complex projective spaces, endowed with their canonical Kähler structure.

The strategy is based on the two remarks, first that the previous condition (1.) implies by [Ren91, lemme 1.3 p.7] that the “classical” groupoid G_0 must be a bundle of commutative groups¹, and second that an example of such isomorphism $\mathcal{I}$ is the Fourier-Gelfand transform $C(\mathbb{T}^n) \simeq C^*(\mathbb{Z}^n)$. This paper is organized as follows:

1. For a $\mathbb{T}^n$ -space M , we construct an isomorphism $C(M) \simeq C^*(G_0)$ for a convenient bundle of commutative groups G_0 , under the assumption that the projection $M \twoheadrightarrow M/\mathbb{T}^n$ has a continuous section.
2. Using a second action on $M/\mathbb{T}^n$ we construct then a deformation groupoid bundle G over $\mathbb{R}$, such that the fiber at 0 of G is G_0 . The sub- $*$ -algebra $\mathcal{Q}$ considered is made of restrictions of C_c^∞ functions on a Lie groupoid $\tilde{G}$ containing G as subgroupoid.
3. We investigate which Poisson structure corresponds to this deformation structure on M .
4. We verify that this construction can be applied to toric manifolds such that the Poisson structure obtained equals the original symplectic structure of the toric manifold.

1 Fourier-Gelfand isomorphisms for a $\mathbb{T}^n$ -space

For G a groupoid, we denote $G^{(0)}$ its units space, and $G \xrightleftharpoons[r]{r} G^{(0)}$ its range and source maps. Recall the definition [Ren]:

Definition 1.1 — A locally compact, Hausdorff, separable groupoid G is *étale* when the fibers of the range map:

$$G^y = \{g \in G \mid r(g) = y\}, \quad y \in G^{(0)},$$

are discrete, and the counting measure is a continuous Haar system.

Let us recall some known properties:

- An open subgroupoid of an étale groupoid is itself – locally compact, Hausdorff, separable – étale.
- The maps r, s are automatically open [Ren]. And the converse is true for bundle of commutative groups: such topological bundle is étale if and only if the projection $G \xrightarrow{r=s} G^{(0)}$ is open (cf. [Ren91]).

¹Remark that a bundle of commutative groups is always amenable. Then replacing globally continuous fields of C^* -algebras by fields only continuous at zero, one can define non globally amenable deformation groupoids

Let M be a Hausdorff, locally compact, separable space endowed with an action α of $\mathbb{T}^n$ denoted by: $(s, x) \in \mathbb{T}^n \times M \longmapsto \alpha_s(x) \in M$. We construct a bundle of commutative groups in the following way. Let us introduce some notations:

- $\Delta = M/\mathbb{T}^n$ denotes the quotient space, which is Hausdorff, separable and locally compact, and $M \xrightarrow{J} \Delta$ is the canonical projection;
- for $y \in \Delta$, $\mathbb{T}_y^n$ is the common isotropy subgroup of all points $x \in M$ such that $J(x) = y$, i.e. $\mathbb{T}_y^n = \{s \in \mathbb{T}^n \mid \alpha_s(x) = x\}$;
- $\mathbb{T}^n/\mathbb{T}_y^n$ is the quotient group and $\widehat{\mathbb{T}^n/\mathbb{T}_y^n}$ is its Pontryagin dual. Since $\mathbb{T}_y^n$ is closed in $\mathbb{T}^n$, the dual map of $\mathbb{T}^n \longrightarrow \mathbb{T}^n/\mathbb{T}_y^n$ is into:

$$\widehat{\mathbb{T}^n/\mathbb{T}_y^n} \hookrightarrow \widehat{\mathbb{T}^n} = \mathbb{Z}^n;$$

- For $k = (k_1, \dots, k_n) \in \mathbb{Z}^n$ and $s = (s_1, \dots, s_n) \in \mathbb{T}^n$, we denote the duality bracket

$$\langle k, s \rangle = s_1^{k_1} \dots s_n^{k_n}.$$

- For every $k \in \mathbb{Z}^n$, we denote: $\Delta(k) = \{y \in \Delta \mid s \in \mathbb{T}_y^n \Rightarrow \langle k, s \rangle = 1\}$. Then $k \in \widehat{\mathbb{T}^n/\mathbb{T}_y^n}$ holds if and only if $y \in \Delta(k)$. We denote moreover: $\Delta(\infty) = \{y \in \Delta \mid \mathbb{T}_y^n = \{1\}\}$.

Finally we set G_0 be the bundle:

$$G_0 = \bigcup_{y \in \Delta} \widehat{\mathbb{T}^n/\mathbb{T}_y^n} = \bigcup_{k \in \mathbb{Z}^n} \Delta(k) \times \{k\} \subset \Delta \times \mathbb{Z}^n,$$

endowed with the topology induced by $\Delta \times \mathbb{Z}^n$.

Proposition 1.2 — *If there exists a continuous section σ of $J : M \xrightarrow{\sigma} \Delta$, then $G_0 \longrightarrow \Delta$ is an étale bundle of commutative groups.*

Proof. — Since the trivial bundle of groups $\Delta \times \mathbb{Z}^n$ is clearly étale, it suffices to prove that G_0 is open. So it suffices to prove that the $\Delta(k)$ are open in Δ , i.e. that their complements $\Delta(k)^c$ are closed in Δ .

First we give a characterization of elements in $\Delta(k)^c$. For each $y \in \Delta$ define a subgroup of $\mathbb{T}$: $H_y = \{\langle k, s \rangle \mid s \in \mathbb{T}_y^n\}$. We have

$$y \in \Delta(k)^c \Leftrightarrow k \notin \widehat{\mathbb{T}^n/\mathbb{T}_y^n} \Leftrightarrow H_y \neq \{1\}.$$

Moreover, recalling that Lie groups do not have small nontrivial subgroups [MZ], in the particular case of a subgroup H of $\mathbb{T}$ we have:

$$(\forall t \in H, |t - 1| < \sqrt{2}) \Leftrightarrow H = \{1\}.$$

Thus

$$y \in \Delta(k)^c \Leftrightarrow (\exists g \in \mathbb{T}_y^n, |\langle k, s \rangle - 1| \geq \sqrt{2}).$$

Second we deduce that for each sequence $(y_i)_i$ in $\Delta(k)^c$ converging to y , then y is itself in $\Delta(k)^c$. Since $y_i \in \Delta(k)^c$, for each i there exists a $s_i \in \mathbb{T}_{y_i}^n$ such that $|\langle k, s_i \rangle - 1| \geq \sqrt{2}$. Let $(s_{i_j})_j$ be a subsequence of $(s_i)_i$ in $\mathbb{T}^n$ converging to a s . Then

$|\langle k, s \rangle - 1| \geq \sqrt{2}$. Moreover, since $s_i \in \mathbb{T}_{y_i}^n$, we have $\alpha_{s_{i_j}}(\sigma(y_{i_j})) = \sigma(y_{i_j})$, for every j . So, taking the limit, we get $\alpha_s(\sigma(y)) = \sigma(y)$, hence $s \in \mathbb{T}_y^n$. Then we have proved :

$$\exists s \in \mathbb{T}_y^n, |\langle k, s \rangle - 1| \geq \sqrt{2}.$$

■

As a corollary, note that the bundle map $G_0 \xrightarrow{p} \Delta$ is open.

Such continuous sections σ may not exist; from now on we suppose that it does and we introduce more notations which may depend implicitly of this σ :

- We define the map $\Delta \times \mathbb{T}^n \xrightarrow{\rho} M$ by: $\rho(y, s) = \alpha_s(\sigma(y))$.
- The “dual” bundle of G_0 is:

$$\widehat{G}_0 = \bigcup_{y \in \Delta} \mathbb{T}^n / \mathbb{T}_y^n,$$

endowed with the biggest topology such that the canonical projection $\Delta \times \mathbb{T}^n \xrightarrow{\widehat{\pi}} \widehat{G}_0$ is continuous.

- Since $\rho(y, s) = \rho(y', s')$ holds if and only if $y = y'$ and $s = s' \bmod \mathbb{T}_y^n$ the map ρ has a one-to-one quotient map $\bar{\rho}$ such that the following diagram is commutative:

$$\begin{array}{ccc} \widehat{G}_0 & \xrightarrow{\bar{\rho}} & M \\ \widehat{\pi} \uparrow & \nearrow \rho & \uparrow \alpha \\ \Delta \times \mathbb{T}^n & \xrightarrow{\sigma \times id_{\mathbb{T}^n}} & M \times \mathbb{T}^n \end{array}$$

- For every $k \in \mathbb{Z}^n \cup \{\infty\}$, we define M_k be the following subset of M :

$$M_k = \rho(\Delta(k) \times \mathbb{T}^n) = J^{-1}(\Delta(k)) = \{\alpha_s(\sigma(y)) \mid s \in \mathbb{T}^n, y \in \Delta(k)\}.$$

In particular, note that $M_\infty = J^{-1}(\Delta(\infty))$ is the maximal stable subset of M where the action of $\mathbb{T}^n$ is free.

- For every $k \in \mathbb{Z}^n$, then define a function k_σ by:

$$\begin{array}{ccc} & M_k & \\ \rho \nearrow & & \searrow k_\sigma \\ \Delta(k) \times \mathbb{T}^n & \xrightarrow{k} & \mathbb{T} \end{array} \quad k_\sigma(\rho(y, s)) = \langle k, s \rangle$$

This is the main result of the section:

Theorem 1.3 — *For every continuous section σ of J and all $k \in \mathbb{Z}^n$, the sets M_k are open in M , and the functions k_σ are continuous. Moreover, for every continuous section σ , there is an isomorphism of C^* -algebras $C_0(M) \xrightarrow[\sim]{\mathcal{I}} C^*(G_0)$, such that, for every $f \in C_c(G_0)$ and $x \in M$:*

$$(\mathcal{I}^{-1}f)(x) = \sum_{k \in \widehat{\mathbb{T}^n / \mathbb{T}_y^n}} f(J(x), k) k_\sigma(x), \quad \text{where } y = J(x).$$

We decompose the proof in several lemmas:

Lemma 1.4 — *The map $\bar{\rho}$ is an homeomorphism.*

Proof. — Since the group $\mathbb{T}^n$ is compact, its action α on M is a proper map, and σ is proper too, since, for every compact subset K of M , the set $\sigma^{-1}(K) = J(K)$ is compact. Then $\bar{\rho} \circ \hat{\pi} = \alpha \circ (\sigma \times id_{\mathbb{T}^n})$ is a closed and continuous map. Then $\bar{\rho}$ itself is closed, since $\hat{\pi}$ is continuous and surjective, and $\bar{\rho}$ is continuous too, by the choice of topology of $\widehat{G}_0$, hence $\bar{\rho}$ is a homeomorphism. ■

Easy consequences of the lemma are:

- the bundle map $\widehat{G}_0 \xrightarrow{\hat{\rho}=J \circ \bar{\rho}} \Delta$ is continuous and proper;
- the bundle duality bracket $\widehat{G}_0 *_\Delta G_0 \longrightarrow \mathbb{T}$ is continuous, denoting $\widehat{G}_0 *_\Delta G_0$ the fibered product.

Then we have:

Lemma 1.5 — *For every function $f \in C^*(G_0)$ let us define its Fourier transform as the function:*

$$\begin{array}{ccc} \widehat{G}_0 & \xrightarrow{\mathcal{F}f} & \mathbb{C} \\ (y, s \bmod \mathbb{T}_y^n) & \mapsto & \sum_{k \in \widehat{\mathbb{T}^n / \mathbb{T}_y^n}} f(y, k) \langle k, s \rangle. \end{array}$$

Then the Fourier transform is an isomorphism of C^ -algebras*

$$C^*(G_0) \xrightarrow{\mathcal{F}} C_0(\widehat{G}_0).$$

Proof. — Since $\widehat{G}_0$ is homeomorphic to M , the space $\widehat{G}_0 \times G_0$ is normal, and $\widehat{G}_0 *_\Delta G_0$ is a closed subset. Then, using Uhrysohn's Theorem, for $f \in C_c(G_0)$ the continuous map $(y, s, k) \longmapsto f(y, k) \langle k, s \rangle$ defined on $\widehat{G}_0 *_\Delta G_0$ admits a continuous extension K on $\widehat{G}_0 \times G_0$. Then using [BouTG, chap X, §3, n° 4, th. 3] and the fact that G_0 is étale hence the counting measure is a *continuous* Haar system, we get the continuity of $\mathcal{F}f$ since:

$$(\mathcal{F}f)(y, s \bmod \mathbb{T}_y^n) = \sum_{k \in \widehat{\mathbb{T}^n / \mathbb{T}_y^n}} K((y, s), (y, k)).$$

Moreover, using

$$|\mathcal{F}f(y, s \bmod \mathbb{T}_y^n)| \leq \sum_{k \in \widehat{\mathbb{T}^n / \mathbb{T}_y^n}} |f(y, k)|,$$

$\mathcal{F}f$ has its support contained in the set $\hat{p}^{-1}(p(\text{Supp } f))$ which is compact, since $\hat{p}$ is proper.

We consider $C_c(G_0)$ as a dense sub- $*$ -algebra of $C^*(G_0)$, and $C_c(\widehat{G}_0)$ as a sub- $*$ -algebra of $C_0(\widehat{G}_0)$, dense too for the sup norm. It is then an easy calculation to verify that the map $C_c(G_0) \xrightarrow{\mathcal{F}} C_c(\widehat{G}_0)$ is a morphism of commutative $*$ -algebras.

Let us prove now, that this $\mathcal{F}$ is isometric. For $f \in C_c(G_0)$ (resp. $\mathbf{f} \in C_c(\widehat{G_0})$) and $y \in \Delta$, we denote f_y (resp. $\mathbf{f}_y$) its restriction to $\widehat{\mathbb{T}^n/\mathbb{T}_y^n}$ (resp. $\mathbb{T}^n/\mathbb{T}_y^n$). Then, from the definition of Fourier transform we have $(\mathcal{F}f)_y = (\mathcal{F}_y)(f_y)$, where $C^*(\widehat{\mathbb{T}^n/\mathbb{T}_y^n}) \xrightarrow{\mathcal{F}_y} C(\mathbb{T}^n/\mathbb{T}_y^n)$ is the classical isometric Fourier-Gelfand transform for commutative groups. Since the bundle map $G_0 \xrightarrow{p} \Delta$ is open we get a continuous field of C^* -algebras $(C^*(\widehat{\mathbb{T}^n/\mathbb{T}_y^n}))_{y \in \Delta}$, which C^* -algebras of continuous sections is $C^*(G_0)$; then, for every $f \in C_c(G_0)$ we get:

$$\|f\|_{C^*(G_0)} = \sup_{y \in \Delta} \|f_y\|_{C^*(\widehat{\mathbb{T}^n/\mathbb{T}_y^n})} = \sup_{y \in \Delta} \|\mathcal{F}_y(f_y)\|_{C(\mathbb{T}^n/\mathbb{T}_y^n)} = \|\mathcal{F}f\|_{C_0(\widehat{G_0})}.$$

Hence, by completion, $\mathcal{F}$ has an isometric extension $C^*(G_0) \xrightarrow{\mathcal{F}} C_0(\widehat{G_0})$. Moreover this extension is surjective, hence an isomorphism, since $\mathcal{F}(C^*(G_0))$ is a closed sub- $*$ -algebra and dense, using the Stone-Weierstrass Theorem, because, for every distinct $(y, s \bmod \mathbb{T}_y^n)$ and $(y', s' \bmod \mathbb{T}_{y'}^n)$ in $\widehat{G_0}$, it is easy to construct a $f \in C_c(G_0)$ such that $(\mathcal{F}f)(y, s \bmod \mathbb{T}_y^n) \neq (\mathcal{F}f)(y', s' \bmod \mathbb{T}_{y'}^n)$. ■

Now we give the proof of the theorem:

Proof. — The sets $M_k = J^{-1}(\Delta(k))$ are open since J is continuous and the $\Delta(k)$ are open. Moreover since, for all $s \in \mathbb{T}^n$ and $y \in \Delta(k)$:

$$k_\sigma(\rho(y, s)) = \langle k, s \rangle,$$

then $k_\sigma = k \circ \bar{\rho}^{-1}$, hence k_σ is continuous.

Then the isomorphism $\mathcal{I}$ is given by:

$$\begin{array}{ccc} C_0(M) & \xrightarrow{\mathcal{I}} & C^*(G_0) \\ \bar{\rho}^* \searrow & & \swarrow \mathcal{F} \\ & C_0(\widehat{G_0}) & \end{array} \quad \mathcal{I}\mathbf{f} = \mathcal{F}^{-1}(\mathbf{f} \circ \bar{\rho}) \Leftrightarrow \mathcal{I}^{-1}f = (\mathcal{F}f) \circ \bar{\rho}^{-1}$$

■

2 Deformation groupoid of a $\mathbb{T}^n$ -space

Let the $\mathbb{T}^n$ -space M is in fact a manifold, and the quotient space Δ is moreover a locally closed subset of a Hausdorff manifold N endowed with an action β of $\mathbb{R}^n$ by diffeomorphism,

$$\begin{array}{ccc} M & \xrightarrow{J} & N \\ \circlearrowleft_\alpha & & \circlearrowleft_\beta \end{array} \quad \Delta \subset N.$$

Then one can construct a deformation groupoid of M (for some Poisson structure) in the following way. For such a given action β there is a right action of $\mathbb{Z}^n$ on $\mathbb{R} \times N$ defined, for every $\hbar \in \mathbb{R}, y \in N, k \in \mathbb{Z}^n$ by :

$$(\hbar, y).k = (\hbar, \beta_{\hbar k}(y)).$$

Let us denote $\tilde{G} = (\mathbb{R} \times N) \rtimes \mathbb{Z}^n$ the cross-product groupoid (cf. [Ren]); then $\tilde{G}$ is an étale Lie bundle of groupoids parametrized by $\hbar \in \mathbb{R}$.

Using the notations $\Delta(k)$ and $\mathbb{T}_y^n$ of the previous section we set:

$$G = \left\{ (\hbar, y, k) \in \mathbb{R} \times N \times \mathbb{Z}^n \mid y, \beta_{\hbar k}(y) \in \Delta(k) \text{ and } \mathbb{T}_y^n = \mathbb{T}_{\beta_{\hbar k}(y)}^n \right\}$$

Then the fiber $\{(\hbar, y, k) \in G \mid \hbar = 0\}$ is exactly the groupoid G_0 defined in the previous section.

Proposition 2.1 — *Let M be a Hausdorff separable manifold with an action of $\mathbb{T}^n$ such that the quotient space Δ is a locally closed part of a Hausdorff manifold N , which is endowed with an action β of $\mathbb{R}^n$ by diffeomorphism. Then the previously defined G is a subgroupoid of $\tilde{G}$, with space of units $G^{(0)} = \mathbb{R} \times \Delta$. Moreover, setting $G_k = G \cap (\mathbb{R} \times N \times \{k\})$, the groupoid G is étale if and only if, for every $k \in \mathbb{Z}^n$, the sets*

$$s(G_k) = \left\{ (\hbar, y) \mid y, \beta_{\hbar k}(y) \in \Delta(k) \text{ and } \mathbb{T}_y^n = \mathbb{T}_{\beta_{\hbar k}(y)}^n \right\}$$

are open in $\mathbb{R} \times \Delta$; then G is a Hausdorff locally compact separable amenable bundle of groupoids, with continuous haar system and open projection $G \longrightarrow \mathbb{R}$.

Proof. — For every groupoid G with space of units $G^{(0)}$, recall that, for every subset U of $G^{(0)}$, the set $r^{-1}(U) \cap s^{-1}(U)$ is a subgroupoid of G , called the *restriction of G to U* , denoted $G|_U$, and which space of units is U .

For every closed subgroup T of $\mathbb{T}^n$, $\widehat{\mathbb{T}^n/T}$ is a subgroup of $\mathbb{Z}^n$, hence $(\mathbb{R} \times N) \rtimes \widehat{\mathbb{T}^n/T}$ a subgroupoid of $\tilde{G}$. Moreover, one can define the set

$$\Delta_T = \{y \in \Delta \mid \mathbb{T}_y^n = T\}.$$

Then we get a partition: $\Delta = \bigcup_T \Delta_T$ and so we can form the restriction

$\left((\mathbb{R} \times N) \rtimes \widehat{\mathbb{T}^n/T} \right) \Big|_{\mathbb{R} \times \Delta_T}$. Since, for every $k \in \mathbb{Z}^n$, we have the partition:

$$\Delta(k) = \bigcup_{T \text{ s.t. } k \in \widehat{\mathbb{T}^n/T}} \Delta_T,$$

then it is easy to verify that G is the disjoint union $G = \bigcup_T \left((\mathbb{R} \times N) \rtimes \widehat{\mathbb{T}^n/T} \right) \Big|_{\mathbb{R} \times \Delta_T}$,

hence a subgroupoid of $\tilde{G}$.

A groupoid with open source and range is étale if and only if it admits a covering by open bisections. Hence a subgroupoid G of an étale groupoid $\tilde{G}$ is itself étale if and only if $G^{(0)}$ is locally closed in $\tilde{G}^{(0)}$, and the images by the source map s (or by the range r) of open bisections covering G are open in $G^{(0)}$. For this particular case, the G_k are bisections covering G .

The bundle map $G \xrightarrow{p} \mathbb{R}$ is open since the first projection $\mathbb{R} \times \Delta \xrightarrow{pr_1} \mathbb{R}$ is open, s is open, and $p = pr_1 \circ s$. Other conditions are particular cases of standard facts for groupoids (cf. [AR]). ■

The main result of the section is

Theorem 2.2 — *Let M be a Hausdorff separable manifold with a continuous action of $\mathbb{T}^n$ such that the quotient space Δ is a locally closed part of a Hausdorff manifold N endowed with an action β of $\mathbb{R}^n$ by diffeomorphisms. If:*

- *the groupoid G is étale (cf. prop. 2.1),*

- the projection $M \xrightarrow{J} N$ is smooth and has a continuous section σ , such that, for every $k \in \mathbb{Z}^n$, the functions $M_k \xrightarrow{k_\sigma} \mathbb{T}$ are smooth,

then M admits a Poisson bracket $\{.,.\}_\Omega$ such that G is a deformation groupoid of (M, Ω) .

This result is closed to the one of Rieffel [Rie] up to quite different technical conditions. The proof consists in making use of Proposition 0.5: we have just seen, in the previous proposition, that G is a convenient groupoid when étale; the condition 1 of Proposition 0.5 comes from Theorem 1.3. For the purpose of condition 2, for every locally closed subset Y of a manifold $\tilde{Y}$, we define the space $C_c^\infty(Y \subset \tilde{Y})$ to be the space of continuous compactly supported functions $f \in C_c(Y)$ which admit a smooth extension $\tilde{f} \in C^\infty(\tilde{Y})$. Then we get the following lemmas.

Lemma 2.3 — *With the assumptions of the Theorem 2.2, then $\mathcal{Q} = C_c^\infty(G \subset \tilde{G})$ is a dense sub- $*$ -algebra of $C^*(G)$.*

Proof. — The space $C_c^\infty(G \subset \tilde{G})$ is clearly a sub-vector space of $C_c(G)$ and is stable by involution, since, for each f with extension $\tilde{f} \in C^\infty(\tilde{G})$, the map f^* has an extension $g \mapsto \tilde{f}(g^{-1})$ which is smooth, since the inverse map of $\tilde{G}$ is a diffeomorphism.

Since the open bisections $G_k = G \cap (\mathbb{R} \times N \times \{k\})$, for $k \in \mathbb{Z}^n$, form a partition of G we get

$$C_c^\infty(G \subset \tilde{G}) = \bigoplus_{k \in \mathbb{Z}^n} C_c^\infty(G_k \subset \mathbb{R} \times N \times \{k\}).$$

Then, to prove the stability of $C_c^\infty(G \subset \tilde{G})$ by product, one has only to show that, for every $l, m \in \mathbb{Z}^n$ and every $f \in C_c^\infty(G_k \subset \mathbb{R} \times N \times \{l\})$, $f' \in C_c^\infty(G_l \subset \mathbb{R} \times N \times \{m\})$, then $f * f'$ is in $C_c^\infty(G \subset \tilde{G})$. This comes easily from the formula:

$$(f * f')(h, y, k) = \begin{cases} f(h, \beta_{hl}(y), l) f'(h, y, m) & \text{if } k = l + m \\ 0 & \text{otherwise.} \end{cases}$$

Since $C_c(G)$ is dense in $C^*(G)$, then $C_c^\infty(G \subset \tilde{G})$ is dense in $C^*(G)$ if every $f \in C_c(G)$ can be approached uniformly by functions $f_n \in C_c^\infty(G \subset \tilde{G})$ since the topology induced by $C^*(G)$ on $C_c(G)$ is the sup-norm topology. To prove this we remark, since G is locally closed in $\tilde{G}$, that a function $f_n \in C_c(G)$ is in $C_c^\infty(G \subset \tilde{G})$ if and only if it admits a smooth and compactly supported extension $\tilde{f}_n \in C_c^\infty(\tilde{G})$. For every $f \in C_c(G)$, since its support is compact, f admits a continuous extension $\tilde{f}$ on $\tilde{G}$, such that $\text{Supp } \tilde{f} \cap G$ is compact. Then this extension $\tilde{f}$ can be uniformly approximated by $\tilde{f}_n \in C_c^\infty(\tilde{G})$; and using smooth partition of unity we can suppose that $\text{Supp } \tilde{f}_n \subset \text{Supp } \tilde{f}$. Then the restrictions of the $\tilde{f}_n$ to G , $f_n = \tilde{f}_n|_G$, are continuous and have their supports included in $\text{Supp } \tilde{f} \cap G$ hence compact; so, using the previous characterization, we have $f_n \in C_c^\infty(G \subset \tilde{G})$ and :

$$\sup_G |f - f_n| \leq \sup_{\tilde{G}} |\tilde{f} - \tilde{f}_n| \xrightarrow{n \rightarrow \infty} 0.$$

■

As a corollary $\mathcal{Q}_0 = C_c^\infty(G_0 \subset \tilde{G}_0)$ is dense in $C^*(G_0)$ (condition 2a). And from the formula:

$$(\mathcal{I}^{-1}f_0)(x) = \sum_{k \in \widehat{\mathbb{T}^n / \mathbb{T}_{J(x)}^n}} f_0(J(x), k) k_\sigma(x),$$

of Theorem 1.3 and the assumptions that J and the k_σ are smooth, we get $\mathcal{I}^{-1}\mathcal{Q}_0 \subset C_0^\infty(M)$ (condition 2b). The condition 2c', hence the Theorem 2.2 follows from the lemma:

Lemma 2.4 — *With the assumptions of the Theorem 2.2, for every $f, f' \in C_c^\infty(G \subset \tilde{G})$ and every $\tilde{e} \in C_c^\infty(\tilde{G})$ which extends $f * f' - f' * f$ on $\tilde{G}$ (resp. on a neighborhood in $\tilde{G}$ of a given point of G_0), then the restriction $d = \tilde{d}|_G$ of $\tilde{d} \in C^\infty(\tilde{G})$ defined by:*

$$\tilde{d}(\hbar, y, k) = \frac{1}{i} \int_0^1 \frac{\partial \tilde{e}}{\partial \hbar}(\hbar t, y, k) dt,$$

1. *is a continuous extension of the section $\hbar \mapsto \frac{1}{i\hbar}[f_\hbar, f'_\hbar]$ on G (resp. on a neighborhood in G of the considered point of G_0).*
2. *has the same support as $f * f' - f' * f$ hence compact*
3. *has value on G_0 (resp. at the considered point of G_0) only depending on f_0 and f'_0 , and given by :*

$$\{f_0, f'_0\}_0(0, y, k) = \frac{1}{i} \frac{\partial \tilde{e}}{\partial \hbar}(0, y, k).$$

Proof. —

1. The 0-order Taylor integral formula gives:

$$\tilde{e}(\hbar, y, k) = \tilde{e}(0, y, k) + i\hbar \tilde{d}(\hbar, y, k).$$

For every $(\hbar, y, k)$ in G (resp. in the considered neighbourhood of G_0), since $C^*(G_0)$ is commutative, and since $(0, y, \hbar)$ is in G_0 , we get:

$$\tilde{e}(0, y, k) = (f * f' - f' * f)(0, y, k) = 0.$$

Hence, on G , we get:

$$f * f' - f' * f = i\hbar d,$$

i.e. d is a continuous extension of $\frac{f * f' - f' * f}{i\hbar}$.

2. Since the bundle map $G \longrightarrow \mathbb{R}$ is open, the set $G - G_0$ is dense in G , hence the continuous extension d is unique, and the previous relation $f * f' - f' * f = i\hbar d$ shows that $f * f' - f' * f$ and d have the same support.
3. Since the map $(f, f') \mapsto f * f' - f' * f$ is bilinear, so is $(f, f') \mapsto d|_{G_0}$ too. Then it is sufficient to prove that $f_0 = 0$ implies $d|_{G_0} = 0$. In the same way as for $\tilde{e}$ (we had $\tilde{e}|_{G_0} = 0$) we get:

$$\exists \delta f \in C_c^\infty(G \subset \tilde{G}), \quad f = i\hbar \delta f.$$

Then $\frac{f * f' - f' * f}{i\hbar} = \delta f * f' - f' * \delta f$; this can be continuously extended by 0 at $\hbar = 0$. ■

3 Computation of the Poisson bracket

With some additional technical assumptions to the conditions of the Theorem 2.2 one can compute explicitly the Poisson structure Ω on M such that G is a deformation groupoid of (M, Ω) . We introduce some notations and conventions. We identify $\mathbb{R}^n$ with its Lie algebra $Lie(\mathbb{R}^n)$. We fix here a basis $E_1, \dots, E_n$ of $\mathbb{R}^n$, such that we identify moreover $\mathbb{R}^n$ with $Lie(\mathbb{T}^n)$, the Lie algebra of $\mathbb{T}^n$, and with its linear dual $Lie(\mathbb{T}^n)^*$; in particular $\mathbb{Z}^n = \widehat{\mathbb{T}^n}$ is viewed as a lattice in $\mathbb{R}^n$, and denoting $(\cdot | \cdot)$ the duality bracket of $\mathbb{R}^n$, and $Lie(\mathbb{T}^n) \xrightarrow{\exp} \mathbb{T}^n$ the exponential map, for every $k = (k_1, \dots, k_n) \in \mathbb{Z}^n$ and every $X \in \mathbb{R}$, we get:

$$(E_i | E_j) = \delta_{i,j}, \quad (k | E_i) = k_i, \quad \langle k, \exp X \rangle = e^{i(k|X)}.$$

Moreover the vector field on M (resp. N) of the infinitesimal action of α (resp. β) in the direction $X \in \mathbb{R}^n$ will be denoted by ξ_X^α (resp. ξ_X^β) i.e. :

$$\forall x \in M, \quad \xi_X^\alpha(x) = \left. \frac{d}{dh} \alpha_{\exp(hX)}(x) \right|_{h=0}, \quad \text{and} \quad \forall y \in N, \quad \xi_X^\beta(y) = \left. \frac{d}{dh} \beta_{hX}(y) \right|_{h=0}.$$

We denote $E_1^*, \dots, E_n^*$ the functions on G_0 defined by:

$$E_i^*(0, y, k) = (k | E_i).$$

The first step is:

Proposition 3.1 — *With the assumptions of Theorem 2.2, let Ω be the Poisson bivector on M such that G is a deformation groupoid of (M, Ω) . Then, for every $f, f' \in C_c^\infty(G \subset \tilde{G})$, every extension $\tilde{f}, \tilde{f}' \in C_c^\infty(\tilde{G})$ respectively of f and f' , denoting $f_0 = f|_{G_0}$ and $f'_0 = f'|_{G_0}$, and $\mathcal{I}$ being the isomorphism provided by Theorem 1.3, we get:*

$$\{ \mathcal{I}^{-1} f_0, \mathcal{I}^{-1} f'_0 \}_\Omega = \frac{1}{i} \sum_{i=1}^n \mathcal{I}^{-1} \left[d\tilde{f}(\xi_{E_i}^\beta) \Big|_{G_0} \right] \left[\mathcal{I}^{-1}(E_i^* f'_0) \right] - \left[\mathcal{I}^{-1}(E_i^* f_0) \right] \mathcal{I}^{-1} \left[d\tilde{f}'(\xi_{E_i}^\beta) \Big|_{G_0} \right].$$

Proof. — Since the Poisson bracket $\{ \cdot, \cdot \}_\Omega$ on M is given by the relation:

$$\mathcal{I}^{-1} \{ f_0, f'_0 \}_0 = \{ \mathcal{I}^{-1} f_0, \mathcal{I}^{-1} f'_0 \}_\Omega,$$

we have only to prove that:

$$\{ f_0, f'_0 \}_0 = \frac{1}{i} \sum_{i=1}^n d\tilde{f}(\xi_{E_i}^\beta) \Big|_{G_0} * (E_i^* f'_0) - (E_i^* f_0) * d\tilde{f}'(\xi_{E_i}^\beta) \Big|_{G_0}.$$

We can restrict to functions f, f' with extensions $\tilde{f}, \tilde{f}'$ both supported in $\mathbb{R} \times N \times \{l\}$ and $\mathbb{R} \times N \times \{m\}$, for some $l, m \in \mathbb{Z}^n$. Then $f * f'$ and $f' * f$ are both supported in $\mathbb{R} \times N \times \{m+l\}$

Since we have supposed that G is étale, then every element in $G_0 \cap (\mathbb{R} \times N \times \{m+l\})$ has a neighborhood V in G such that, for every $(h, y, m+l)$ in V , we have together:

$$(h, y, l), (h, \beta_{hl}(y), m), (h, y, m), (h, \beta_{hm}(y), l) \in G.$$

Then, on V , we get:

$$f * f' - f' * f = \tilde{e},$$

where $\tilde{e} \in C^\infty(\tilde{G})$ is the function defined by:

$$\tilde{e}(\hbar, y, m + l) = \tilde{f}(\hbar, \beta_{hl}(y), m) \tilde{f}'(\hbar, y, l) - \tilde{f}'(\hbar, \beta_{hm}(y), l) \tilde{f}(\hbar, y, m).$$

Due to lemma 2.4 we get:

$$\{f_0, f'_0\}_0(0, y, k) = \frac{1}{i} \frac{\partial \tilde{e}}{\partial \hbar}(0, y, k).$$

We compute:

$$\begin{aligned} \frac{\partial \tilde{e}}{\partial \hbar}(\hbar, y, k) &= \left(\frac{\partial \tilde{f}}{\partial \hbar}(\hbar, \beta_{hl}(y), m) + d\tilde{f}\left(\frac{d\beta_{hl}}{d\hbar}(y)\right)(\hbar, \beta_{hl}(y), m) \right) \tilde{f}'(\hbar, y, l) \\ &\quad + \tilde{f}(\hbar, \beta_{hl}(y), m) \frac{\partial \tilde{f}'}{\partial \hbar}(\hbar, y, l) - \tilde{f}'(\hbar, \beta_{hm}(y), l) \frac{\partial \tilde{f}}{\partial \hbar}(\hbar, y, m) \\ &\quad - \left(\frac{\partial \tilde{f}'}{\partial \hbar}(\hbar, \beta_{hm}(y), l) + d\tilde{f}'\left(\frac{d\beta_{hm}}{d\hbar}(y)\right)(\hbar, \beta_{hm}(y), l) \right) \tilde{f}(\hbar, y, m). \end{aligned}$$

For $\hbar = 0$, since $\beta_0(y) = y$ there remains :

$$\frac{\partial \tilde{e}}{\partial \hbar}(0, y, k) = d\tilde{f}(\xi_l^\beta)(0, y, m) \tilde{f}'(0, y, l) - d\tilde{f}'(\xi_m^\beta)(0, y, l) \tilde{f}(0, y, m).$$

Since $\xi_l^\beta = \sum_{i=1}^n (l | E_i) \xi_{E_i}^\beta$ we get then:

$$\{f_0, f'_0\}_0(0, y, k) = \frac{1}{i} \sum_{i=1}^n d\tilde{f}(\xi_{E_i}^\beta)(0, y, m) E_i^*(0, y, l) \tilde{f}'(0, y, l) - d\tilde{f}'(\xi_{E_i}^\beta)(0, y, l) E_i^*(0, y, m) \tilde{f}(0, y, m),$$

i.e.

$$\{f_0, f'_0\}_0 = \frac{1}{i} \sum_{i=1}^n d\tilde{f}(\xi_{E_i}^\beta) \Big|_{G_0} * (E_i^* f'_0) - (E_i^* f_0) * d\tilde{f}'(\xi_{E_i}^\beta) \Big|_{G_0}.$$

■

The second step is to compute $\mathcal{I}^{-1} d\tilde{f}(\xi_{E_i}^\beta) \Big|_{G_0}$ and $\mathcal{I}^{-1}(E_i^* f_0)$ with respect to $\mathcal{I}^{-1} f_0$. We use some additional conditions to obtain:

Theorem 3.2 — *With the assumptions of Theorem 2.2 and suppose moreover that:*

1. *the torus $\mathbb{T}^n$ acts (by α) on M by diffeomorphisms;*
2. *$\Delta(\infty) = \{y \in \Delta \mid \mathbb{T}_y^n = \{1\}\}$ is open in N and dense in Δ – for density it occurs for example when the action α is effective;*
3. *the restricted map $\Delta(\infty) \times \mathbb{T}^n \xrightarrow{\rho} M_\infty$ (cf. lemma 1.4) is a diffeomorphism*

Then the Poisson structure of Theorem 2.2 is given on M_∞ by the bivector:

$$\Omega = \sum_{i=1}^n \xi_{E_i}^\alpha \wedge \rho_*(\xi_{E_i}^\beta).$$

Proof. — Remark that, for $\mathcal{Q} = C_c^\infty(G \subset \tilde{G})$, we have both:

$$\mathcal{Q}_0 = C_c^\infty(G_0 \subset \tilde{G}_0) \text{ and } \mathcal{I}^{-1}\mathcal{Q}_0 \subset C_0^\infty(M),$$

hence $\mathcal{Q}_0$ is dense in $C_0^\infty(M)$ since $C_c^\infty(G_0 \subset \tilde{G}_0)$ is dense in $C^*(G_0)$. So we have only to prove that, for every $f_0, f'_0 \in C_c^\infty(G_0 \subset \tilde{G}_0)$:

$$\begin{aligned} \{ \mathcal{I}^{-1}f_0, \mathcal{I}^{-1}f'_0 \}_\Omega &= \langle d(\mathcal{I}^{-1}f_0) \otimes d(\mathcal{I}^{-1}f'_0) | \Omega \rangle \\ &= \sum_{i=1}^n d(\mathcal{I}^{-1}f_0) \left(\xi_{E_i}^\alpha \right) d(\mathcal{I}^{-1}f'_0) \left(\rho^* \left(\xi_{E_i}^\beta \right) \right) - d(\mathcal{I}^{-1}f'_0) \left(\xi_{E_i}^\alpha \right) d(\mathcal{I}^{-1}f_0) \left(\rho^* \left(\xi_{E_i}^\beta \right) \right). \end{aligned}$$

Moreover, it can be proved only on $M_\infty = J^{-1}(\Delta(\infty))$ since it is dense in M . In fact it is a consequence of both:

$$\mathcal{I}^{-1}(E_i^* f_0) = \frac{1}{\iota} d(\mathcal{I}^{-1}f_0)(\xi_{E_i}^\alpha) \quad \text{and} \quad \mathcal{I}^{-1} d\tilde{f} \left(\xi_{E_i}^\beta \right) \Big|_{G_0} = d(\mathcal{I}^{-1}f_0) \left(\rho_*(\xi_{E_i}^\beta) \right).$$

For the first formula, recall from Theorem 1.3 that, for all $x \in M$:

$$\mathcal{I}^{-1}(f_0)(x) = \sum_{k \in \mathbb{Z}^n} f(0, J(x), k) k_\sigma(x).$$

Using condition (1), since, for all $x \in M$:

$$dk_\sigma(\xi_{E_i}^\alpha)(x) = \frac{d}{dt} k_\sigma \left(\alpha_{\exp(tE_i)}(x) \right) \Big|_{t=0} = \frac{d}{dt} \langle k, \exp(tE_i) \rangle \Big|_{t=0} k_\sigma(x) = \iota(k|E_i) k_\sigma(x),$$

and since J is invariant with respect to the action α , we get, for all $x \in M$:

$$\begin{aligned} d_x(\mathcal{I}^{-1}f_0)(\xi_{E_i}^\alpha) &= \sum_{k \in \mathbb{Z}^n} f(0, J(x), k) dk_\sigma(\xi_{E_i}^\alpha)(x) \\ &= \iota \sum_{k \in \mathbb{Z}^n} f(0, J(x), k) E_i^*(0, J(x), k) k_\sigma(x) = \iota \mathcal{I}^{-1}(E_i^* f_0)(x). \end{aligned}$$

For the second formula, since $\xi_{E_i}^\beta$ is a vector field on $\Delta(\infty)$, using conditions (2) and (3), for all $s \in \mathbb{T}^n$ and $y \in \Delta(\infty)$, we get:

$$\begin{aligned} d_{\rho(s,y)}(\mathcal{I}^{-1}f_0)(\rho_*(\xi_{E_i}^\beta)) &= d_{(s,y)}(\mathcal{I}^{-1}f_0 \circ \rho)(\xi_{E_i}^\beta) \\ &= \sum_{k \in \mathbb{Z}^n} d_y \tilde{f}(\xi_{E_i}^\beta)(k, s) \\ &= \left(\left(\mathcal{I}^{-1} d\tilde{f}(\xi_{E_i}^\beta) \Big|_{G_0} \right) \circ \rho \right)(s, y) \end{aligned}$$

hence: $d(\mathcal{I}^{-1}f_0)(\rho_*(\xi_{E_i}^\beta)) = \mathcal{I}^{-1} \left(d\tilde{f}(\xi_{E_i}^\beta) \Big|_{G_0} \right)$. ■

4 Application to toric manifolds

First we recall some facts concerning toric manifolds. We will use the notations and conventions of sections 1 and 3.

Definition 4.1 — The action α of a Lie group G on a symplectic manifold (M, ω) by diffeomorphism is *Hamiltonian* when there exists a so-called *moment map* $M \xrightarrow{J} \text{Lie}(G)^*$ such that:

$$\forall f \in C^\infty(M), \forall X \in \text{Lie}(G), df(\xi_X^\alpha) = \{ (J|X), f \}_\omega.$$

When M is connected, J is unique up to a constant.

A *toric manifold* is a compact and connected symplectic manifold (M, ω) endowed with an effective hamiltonian action α of $\mathbb{T}^n$ such that M has (real) dimension $2n$.

In the '80s Atiyah [Ati], Guillemin and Sternberg [GS82] proved that, for a toric manifold, the image of the moment map $J(M)$ is a convex polytope in $\mathbb{R}^n = Lie(\mathbb{T}^n)^*$. And Delzant [Del88] completed this result by the following ones:

- the map J is $\mathbb{T}^n$ -invariant and the quotient map is a bijection $M/\mathbb{T}^n \simeq J(M)$, hence an homeomorphism. So, from now on, we identify $\Delta = M/\mathbb{T}^n$ with the polytope $J(M)$ in $N = \mathbb{R}^n$.
- Every isotropy group $\mathbb{T}_y^n$ is connected and depends only on the open face of the polytope Δ containing y ; more precisely there exists a parametrization of Δ such that:

$$\Delta = \{y \in \mathbb{R}^n \mid \forall j \in \{1, \dots, n_0\}, (y \mid X_j) \geq \lambda_j\}, \quad X_j \in \mathbb{Z}^n, \quad \lambda_j \in \mathbb{R}$$

$$Lie(\mathbb{T}_y^n) = span\{X_j \mid (y \mid X_j) = \lambda_j\}.$$

In particular $\Delta(\infty)$ is the topological interior of Δ in $\mathbb{R}^n$.

- Two toric manifolds having the same polytope Δ are diffeomorphic, in a $\mathbb{T}^n$ -equivariant symplectic way. So a toric manifold is totally characterized by its polytope.
- When Δ is the polytope of a toric manifold M , then there is an explicit construction of M (with action an symplectic form ω) from Δ .

We need to recall briefly the main steps of the latter construction; we use the previous parametrization of Δ :

1. $\mathbb{T}^{n_0}$ has a (diagonal) hamiltonian action on $\mathbb{C}^{n_0}$ endowed with its canonical symplectic structure $\omega_0 = \sum_{j=1}^{n_0} dx_j \wedge dy_j$. The moment map is given by:

$$J_0(z_1, \dots, z_{n_0}) = (\lambda_1, \dots, \lambda_{n_0}) + \frac{1}{2}(|z_1|^2, \dots, |z_{n_0}|^2) \in \mathbb{R}^{n_0}.$$

2. There exists a group exact sequence:

$$1 \longrightarrow \mathbb{T}^d \xrightarrow{i} \mathbb{T}^{n_0} \xrightarrow{\pi} \mathbb{T}^n \longrightarrow 1, \quad (n_0 = n + d)$$

such that $d\pi(F_j) = X_j$, where $F_1, \dots, F_{n_0}$ is the canonical basis of $\mathbb{R}^{n_0}$.

3. Using $Lie(\mathbb{T}^p)^* = \mathbb{R}^p$, let $\mathbb{R}^{n_0} \xrightarrow{di^*} \mathbb{R}^d$ be the cotangent map of i , and let us define $J_d = di^* \circ J_0$. Then we have an inclusion map $J_d^{-1}\{0\} \xrightarrow{j} \mathbb{C}^{n_0}$ and $J_d^{-1}\{0\}$ in a stable part of $\mathbb{C}^{n_0}$ for the actions of $\mathbb{T}^{n_0}$ and of $\mathbb{T}^d$ – identified with its image by i in $\mathbb{T}^{n_0}$. Moreover these two actions commute and we get a quotient action of $\mathbb{T}^n \simeq \mathbb{T}^{n_0}/\mathbb{T}^d$ on $J_d^{-1}\{0\}/\mathbb{T}^d$.

4. Using Marsden-Weinstein's symplectic reduction [MW74] the canonical projection $J_d^{-1}\{0\} \xrightarrow{pr} J_d^{-1}\{0\}/\mathbb{T}^d$ is a submersion, hence $J_d^{-1}\{0\}/\mathbb{T}^d$ is a manifold of dimension $2n$, with symplectic structure ω such that $pr^*\omega = j^*\omega_0$. Moreover the action of $\mathbb{T}^n$ is hamiltonian with moment map J . Calculations prove that the image of J is then Δ so we get the following commutative diagram:

$$\begin{array}{ccccc}
 M \simeq J_d^{-1}\{0\}/\mathbb{T}^d & \xleftarrow{pr} & J_d^{-1}\{0\} & \xrightarrow{j} & \mathbb{C}^{n_0} \\
 \searrow J & & \searrow J_0 \circ j & \nearrow J_0 & \searrow J_d \\
 \mathbb{R}^n & \xrightarrow{d\pi^*} & \mathbb{R}^{n_0} & \xrightarrow{di^*} & \mathbb{R}^d
 \end{array}
 \quad
 \begin{array}{l}
 J_0 \circ j = d\pi^* \circ J \circ pr \\
 di^* \circ d\pi^* = 0
 \end{array}$$

The main result of this article is:

Theorem 4.2 — *Let (M, ω) be a toric manifold with the action α of $\mathbb{T}^n$, let $M \xrightarrow{J} N = \mathbb{R}^n$ be the moment map, and let β be the action of $\mathbb{R}^n$ on N by translation:*

$$\beta_X(y) = y + X.$$

Then, with the same notations, the groupoid G of Theorem 2.2 is a deformation groupoid of (M, ω) .

We will just prove that the conditions of Theorem 2.2 are fulfilled and then use Theorem 3.2 to prove that the Poisson structure Ω so obtained is the same as the one coming from ω , i.e. $\{.,.\}_\omega = \{.,.\}_\Omega$.

Lemma 4.3 — *With respect to the assumptions of Theorem 4.2, for every $k \in \mathbb{Z}^n$, the following set is open in $\Delta \times \mathbb{R}^n$:*

$$s(G_k) = \{ (\hbar, k) \in \mathbb{R} \times \Delta \mid y, y + \hbar k \in \Delta(k) \text{ and } \mathbb{T}_y^n = \mathbb{T}_{y+\hbar k}^n \}.$$

Proof. — Recall that the isotropy subgroups $\mathbb{T}_y^n$ are connected and

$$Lie(\mathbb{T}_y^n) = \text{span} \{ X_j \mid (y \mid X_j) = \lambda_j \}.$$

So we get:

$$\Delta(k) = \{ y \in \Delta \mid \forall j \in \{1, \dots, n_0\}, (y \mid X_j) = \lambda_j \Rightarrow (k \mid X_j) = 0 \} = \Delta \cap U(k),$$

where $U(k)$ is the open subset of $\mathbb{R}^n$ defined by:

$$U(k) = \{ y \in \mathbb{R}^n \mid \forall j \in \{1, \dots, n_0\}, (k \mid X_j) \neq 0 \Rightarrow (y \mid X_j) > \lambda_j \}.$$

Since we have

$$(y + \hbar k \mid X_j) = (y \mid X_j) + \hbar (k \mid X_j),$$

then we get that:

$$y \in \Delta(k) \Rightarrow Lie(\mathbb{T}_y^n) \subset Lie(\mathbb{T}_{y+\hbar k}^n).$$

In the same way we get: $y + \hbar k \in \Delta(k) \Rightarrow \text{Lie}(\mathbb{T}_{y+\hbar k}^n) \subset \text{Lie}(\mathbb{T}_y^n)$.
Hence $y, y + \hbar k \in \Delta(k)$ implies that $\mathbb{T}_y^n = \mathbb{T}_{y+\hbar k}^n$ and we have furthermore:

$$y, y + \hbar k \in \Delta(k) \Leftrightarrow y \in \delta(k) \text{ and } y + \hbar k \in U(k).$$

So

$$s(G_k) = \{ (\hbar, k) \in \mathbb{R} \times \Delta \mid y \in \Delta(k) \text{ and } y + \hbar k \in U(k) \}$$

is open in $\mathbb{R} \times \Delta$. ■

For the action of $\mathbb{T}^{n_0}$ in $\mathbb{C}^{n_0}$ we have $J_0(\mathbb{C}^{n_0}) = \lambda + (\mathbb{R}_+)^{n_0} = \Delta_0$, and there is a natural section σ_0 of J_0 given, for every $(w_1, \dots, w_{n_0}) \in \lambda + (\mathbb{R}_+)^{n_0}$ by:

$$\sigma_0(w_1, \dots, w_{n_0}) = \left(\sqrt{2w_1 - \lambda_1}, \dots, \sqrt{2w_{n_0} - \lambda_{n_0}} \right).$$

Lemma 4.4 — *For every toric manifold (M, ω) constructed as before, the map $\sigma = pr \circ \sigma_0 \circ d\pi^*$ is well defined and is a section of $M \xrightarrow{J} \Delta \subset \mathbb{R}^n$.*

Proof. — Using the previous big commuting diagram and $J_0 \circ \sigma_0 = \text{id}$, for every $y \in \Delta$ we get:

$$J_d((\sigma_0 \circ d\pi^*)(y)) = (di^* \circ d\pi^*)(y) = 0,$$

hence $(\sigma_0 \circ d\pi^*)(y)$ is in $J_d^{-1}\{0\}$, so σ is well defined.

The same kind of calculus proves:

$$d\pi^* \circ (J \circ \sigma) = d\pi^*.$$

Since π is surjective, $d\pi^*$ is into, hence $J \circ \sigma = \text{id}$. ■

In particular this section σ is continuous. Moreover, since the restriction of σ_0 to $\Delta_0(0) = \lambda + (\mathbb{R}_+^*)^{n_0}$ is smooth, we get that σ is smooth on $\Delta(\infty)$, hence the map $\Delta(\infty) \times \mathbb{T}^n \xrightarrow{\rho} M$ is smooth.

Lemma 4.5 — *For every toric manifold (M, ω) constructed as before and with respect to the previous σ , for all $k \in \mathbb{Z}^n$, the map $M_k \xrightarrow{k_\sigma} \mathbb{T}$ is smooth.*

Proof. — In the same way as for the action α of $\mathbb{T}^n$ on M and the section σ , considering the diagonal action of $\mathbb{T}^{n_0}$ on $\mathbb{C}^{n_0}$ and the section σ_0 , one can define : a map $\Delta_0 \times \mathbb{T}^{n_0} \xrightarrow{\rho_0} \mathbb{C}^{n_0}$, for every $l = (l_1, \dots, l_{n_0}) \in \mathbb{Z}^{n_0}$ the sets $\Delta_0(l)$ and $\mathbb{C}_l^{n_0} = \rho_0(\Delta_0(l) \times \mathbb{T}^{n_0})$, and finally the map $\mathbb{C}_l^{n_0} \xrightarrow{l_{\sigma_0}} \mathbb{T}$. Explicitely we get:

$$\mathbb{C}_l^{n_0} = \{ (z_1, \dots, z_{n_0}) \in \mathbb{C}^{n_0} \mid l_j \neq 0 \Rightarrow z_j \neq 0 \}$$

and, since $\sigma_0(J_0(z_1, \dots, z_{n_0})) = (|z_1|, \dots, |z_{n_0}|)$:

$$l_{\sigma_0}(z_1, \dots, z_{n_0}) = \prod_{j|l_j \neq 0} \left(\frac{z_j}{|z_j|} \right)^{l_j},$$

hence l_{σ_0} is smooth on $\mathbb{C}_l^{n_0}$.

For a $k \in \mathbb{Z}^n$, we apply this construction to the particular case of $l = d\pi^*(k) \in \mathbb{Z}^{n_0}$. Then one can check that the following diagram is commutative:

$$\begin{array}{ccc} M_k & \xleftarrow{pr} pr^{-1}(M_k) & \xrightarrow{j} \mathbb{C}_l^{n_0} \\ & \searrow k_\sigma & \swarrow l_{\sigma_0} \\ & \mathbb{T} & \end{array} \quad l_{\sigma_0} \circ j = k_\sigma \circ pr$$

Since pr is a submersion and $l_{\sigma_0} \circ j$ is smooth, then k_σ is smooth too. $\blacksquare$

So the conditions of Theorem 2.2 are true. To use Theorem 3.2 there remains only to prove that the map $\Delta(\infty) \times \mathbb{T}^n \xrightarrow{\rho} M_\infty$ is a diffeomorphism; we already know that it is one-to-one and smooth. It comes from:

Lemma 4.6 — *For every toric manifold (M, ω) constructed as before, and every $(y, s) \in \Delta(\infty) \times \mathbb{T}^n$, identifying the tangent bundle of $\Delta(\infty) \times \mathbb{T}^n$ at (y, s) with $\mathbb{R}^n \times \mathbb{R}^n$, we get:*

$$\forall (Y, X), (Y', X') \in \mathbb{R}^n \times \mathbb{R}^n, \quad \rho^* \omega((Y, X), (Y', X')) = (X|Y') - (Y|X').$$

Proof. — For every $y \in \Delta(\infty)$ and $s \in \mathbb{T}^n$ we get:

$$d_{y,s} \rho(Y, X) = \xi_X^\alpha(\rho(y, s)) + d_{\sigma(y)} \alpha_s(d_y \sigma(Y)).$$

So, using the relation between ω and J given by $\omega(\xi_X^\alpha, T) = d(J|X)(T)$ we get:

$$\rho^* \omega((0, X), (0, X')) = \omega(\xi_X^\alpha, \xi_{X'}^\alpha) = 0.$$

$$\rho^* \omega((0, X), (Y, 0)) = \omega(\xi_X^\alpha, d\alpha_s(d\sigma(Y))) = (X|Y).$$

Then after having computed that $\sigma_0^* \omega_0 = 0$, using $pr^* \omega = j^* \omega_0$ we get:

$$\sigma^* \omega = (pr \circ \sigma_0 \circ d\pi^*)^* \omega = (\sigma_0 \circ d\pi^*)^* (pr^* \omega) = (d\pi^*)^* (\sigma_0^* \omega_0) = 0.$$

Hence:

$$\rho^* \omega((Y, 0), (Y', 0)) = (\alpha_s \circ \sigma)^* \omega((Y, 0), (Y', 0)) = \sigma^* \omega((Y, 0), (Y', 0)) = 0.$$

$\blacksquare$

Hence ρ is a diffeomorphism since it pulls back a nondegenerated 2-form on a nondegenerated 2-form.

Moreover, using $\xi_Y^\beta = Y$ since β acts by translations, we get

$$d_{\sigma(y)} \alpha_s(d_y \sigma(Y)) = \rho^* \left(\xi_Y^\beta \right)$$

Hence, it follows immediatly that the Poisson bivector associated to ω equals the bivector Ω given in Theorem 3.2. So Theorem 4.2 is proved.

Finally let us add a complementary result on the structure of the C^* -algebras $C^*(G_\hbar)$ which occur in the deformation:

Proposition 4.7 — *For every $\hbar \in \mathbb{R}$, every irreducible representation of $C^*(G_\hbar)$ has finite dimension. In particular $C^*(G_\hbar)$ is a C^* -algebra of type I.*

Proof. — For $\hbar = 0$, $C^*(G_0)$ is commutative, hence all representations have dimension 1.

For $\hbar \neq 0$, we remark that the action of $G_\hbar$ on its base $G_\hbar^{(0)}$ has no isotropy, i.e. $G_\hbar$ is a so-called principal groupoid. Moreover every orbit of this action is finite hence closed. But every irreducible representation of the C^* -algebra of a principal groupoid is supported by a closed orbit. $\blacksquare$

Acknowledgments : The author is indebted to J. Renault, G. Skandalis and J. Bellissard for very helpful discussions and comments.

References

- [AR] C. ANANTHARAMAN-DELAROCHE AND J. RENAULT. *Amenable groupoids*. Monographie de l'Enseignement Mathématique (Genève), 2000.
- [ATI] M. F. ATIYAH. Convexity and commuting Hamiltonians. *Bull. London Math. Soc.*, 14:1–15, 1982.
- [BFFLS] F. BAYEN, M. FLATO, C. FRONSDAL, A. LICHNEROWICZ, AND D. STERNHEIMER. Deformation theory and quantization i, ii. *Ann. Physics*, 110:61–110, 111–151, 1978.
- [BOU71] N. BOURBAKI. *Topologie Générale*, volume chap. 1-10. Hermann, 1971.
- [BV] J. BELLISSARD AND M. VITTOT. Heisenberg picture and non-commutative geometry of the semi-classical limit in quantum mechanics. *Ann. I.H.P., Physique théorique*, 52(3):175–235, 1990.
- [CAD] F. CADET. *Déformation et quantification par groupoïde des variétés toriques*. Ph.D. Thesis, Université d'Orléans, 2001. <http://www.math.jussieu.fr/~cadet>
- [CDW] A. COSTE, P. DAZORD, AND A. WEINSTEIN. Groupoïdes symplectiques. In *Publications du Département de Mathématiques. Nouvelle Série. A, Vol. 2*, pages i–ii, 1–62. Univ. Claude-Bernard, Lyon, 1987.
- [CL01] A. CONNES AND G. LANDI. Noncommutative manifolds, the instanton algebra and isospectral deformations. *Commun. Math. Phys.*, 221: 141–159, 2001.
- [Co] A. CONNES. *Noncommutative geometry*. Academic Press Inc., San Diego, CA, 1994.
- [DEL88] T. DELZANT. Hamiltoniens périodiques et images convexes de l'application moment. *Bull. Soc. math. France*, 116:315–339, 1988.
- [DF] R.S. DORAN AND J.M.G. FELL. *Representations of $*$ -Algebras, Locally Compact Groups, and Banach $*$ -Algebraic Bundles*, volume 1 (Basic Representation Theory of Groups and Algebras). Academic Press, 1988.
- [DS] P. DAZORD AND D. SONDAZ. Algébroïdes de Lie et variétés de Poisson. Publication du Département de Mathématiques de l'Université de Lyon-I, 1988. nouvelle série, vol. 1/B.
- [GS82] V. GUILLEMIN AND S. STERNBERG. Convexity properties of the moment mapping. *Invent. Math.*, 67:491–513, 1982.
- [LAN98] N.P. LANDSMAN. *Mathematical topics between classical and quantum mechanics*. Springer-Verlag, New York, 1998.

- [LR99] N.P. LANDSMAN AND B. RAMAZAN. *Quantization of Poisson algebras associated to Lie algebroids*, volume Proceedings of the Conference on Groupoids in Physics, Analysis and Geometry (Boulder 20-24 Juin 1999). A.M.S., 1999. Arxiv math-ph/0001005.
- [MW74] J. MARSDEN AND A. WEINSTEIN. Reduction of symplectic manifolds with symmetry. *Rep. Mathematical Phys.*, 5(1):121–130, 1974.
- [MZ] D. MONTGOMERY AND L. ZIPPIN. *Topological Transformation groups*. Wiley, New-York, 1955.
- [RAM98] B. RAMAZAN. *Quantification par Déformation des Variétés de Lie-Poisson*. Ph.D. Thesis, Université d’Orléans, 1998.
- [REN] J. RENAULT. *A Goupoid Approach to C^* -algebras*, volume 793 of *Lecture Notes in Math*. Springer-Verlag, 1980.
- [REN82] J. RENAULT. C^* -algebras of groupoids and foliations. *Proc. Sympos. Pure Math.*, 38:pages 339–350, 1982.
- [REN87] J. RENAULT. Représentation des produits croisés d’algèbres de groupoïdes. *J. Operator Theory*, 18:67–97, 1987.
- [REN91] J. RENAULT. The ideal structure of groupoid crossed-product C^* -algebras. *J. Operator Theory*, 25:3–36, 1991.
- [RIE89] M. A. RIEFFEL. Déformation quantization of Heisenberg manifolds. *Comm. Math. Phys.*, 122:531–562, 1989.
- [RIE] M. A. RIEFFEL. *Déformation Quantization for Actions of $\mathbb{R}^d$* , volume 106. Monography of the A.M.S., 1993.
- [RIE98] M. A. RIEFFEL. Questions on quantization. In *Operator algebras and operator theory (Shanghai, 1997)*, pages 315–326. Amer. Math. Soc., Providence, RI, 1998.
- [VAR01] J. C. VÁRILLY. Quantum symmetry groups of noncommutative spheres. *Comm. Math. Phys.*, 221: 511–523, 2001.